First record of Phrygilus fruticeti coracinus in Peru

During the wet seasons of 2006 (29 November-1 December) and 2008 (24 November–13 December) between 06h00 to 08h00 on sunny days, we recorded lone male Mourning Sierra Finches Phrygilus fruticeti of the race coracinus at two localities, Tripartito (17°29'S 69°28'W; 4,094 m) and Ayro (17°34'S 69°37'W; 4,243 m), in dpto. Tacna, Peru. Both localities lie close to the junction of the Peru / Chile / Bolivia borders. The sierra finches were observed flocking with Bright-rumped Yellow Finch Sicalis uropygialis and Black-hooded Sierra Finch Phrygilus atriceps. We documented our observations with photographs (Fig. 1). Both sites are dominated by typical highland *pajonal* or grassland (Stipa spp.) and tolar (Parastrephia sp.) vegetation, but most of the finches we recorded were close to settlements. Further surveys were made monthly, in January-September 2009, but no more P. f. coracinus were recorded.

Our observations are the first confirmed record of *P. f. coracinus* in Peru and the northernmost record of the subspecies. In addition they also confirm the presence in Peru of two of the three subspecies of *P. fruticeti*, although *P. f. peruvianus* has been considered a weakly defined race and *coracinus* exhibits some intergradation with nominate *fruticeti*¹. Previously, *P. f. coracinus* was known from the



Figure I. Mourning Sierra Finch Phrygilus fruticeti coracinus, Ayro, dpto. Tacna, Peru, 13 December 2008 (Nataly Hidalgo-Aranzamendi)

altiplano of Bolivia west from Lake Poopó, Oruro, and Uyuni, Potosí, to the Chilean border (Sajama and Pampa Olliaga in Oruro, Sacaya in Tarapacá), and had also been recorded at Isluga and Visviri in northern Chile¹⁻³. No other subspecies of *P. fruticeti* permanently inhabit this area, which might indicate that their presence is due to the seasonal availability of some resource, but further evidence is required. Fjeldså¹ suggested that, in some localities, the species could be only a visitor from higher altitudes, which would suggest migratory behaviour. Other Emberizidae were recorded regularly in our study area such as Sicalis uropygialis, Phrygilus atriceps, Ash-breasted Sierra Finch P. plebeius and White-winged Diuca Finch Diuca speculifera. Unexpectedly, the majority of our observations were made around habitation. Typically P. f. coracinus inhabits bushy or rocky slopes with some brush, in arid and semi-arid regions, or pumice slopes with Polylepis shrub at much higher elevations, rarely in open grasslands unless bordered by shrubby vegetation^{1,2}. Our observations might further indicate that the species undertakes as yet undocumented seasonal movements and that the birds we recorded were opportunistically exploiting regional (seasonal) resources.

Acknowledgements

We thank Dr Evaristo Lopez, Director of Museo de Historia Natural de la Universidad Nacional de San Agustín de Arequipa and of the project 'Distribución, hábitat potencial v realizado, v presión de cacería de Chaetophractus nationi (Xenarthra) en el sur del Perú'. which provided logistical support to the 2006 survey; whilst the project 'Conservación de flora y fauna amenazada en la región Tacna', Gobierno Regional Tacna, provided similar support in 2008. K. Kerr improved the English, and José Velasquez, Gustavo Jara, Roberto Quispe, Lizette Tejada, Shirley Huancollo, Elisban Chino

and Romina Ventura aided the field work.

References

- Fjeldså, J. (1993) Phrygilus coracinus Sclater, 1891, is a valid taxon. Bull Brit. Orn. Club 113: 121–126.
- Fjeldså, J. & Krabbe, N. (1990) Birds of the high Andes. Copenhagen: Zool. Mus., Univ. of Copenhagen & Svendborg: Apollo Books.
- Jaramillo, A. (2003) Birds of Chile. Princeton, NJ: Princeton University Press.

L. Mauricio Ugarte and Nataly Hidalgo-Aranzamendi

Museo de Historia Natural de la Universidad Nacional de San Agustín de Arequipa (Scientific Collection, Ornithology Department), Luna Pizarro 925, Los Pinos, Vallecito—Cercado, Arequipa, Peru. E-mails: mugartelewis@yahoo.com and nhidalgoa@hotmail.com.

Received 10 May 2009; final revision accepted 2 December 2009 (published online 16 March 2010)