Cotinga 32 Short Communications



Figure I. Red-crested Finch Coryphospingus cucullatus, Amazonia Lodge, dpto. Madre de Dios, Peru, November 2009 (Virgilio Yábar Calderón)

south-east Peru in tropical lowland evergreen forest.

In Bolivia, Red-crested Finch occurs in dptos. Beni, La Paz, Cochabamba, Santa Cruz, Chuquisaca and Tarija, in habitats including secondary forest and tropical lowland evergreen forest, and the species is considered to be an austral migrant¹. In Brazil it is known from the south and north-east of the country², where the species is occasionally observed at the borders of humid *terra firme* forest (G. M. Kirwan pers. obs.).

Acknowledgements

Barry Walker and Guy Kirwan provided comments on the manuscript.

References

- Hennessey, A. B., Herzog, S. K. & Sagot, F. (2003)
 Lista anotada de las aves de Bolivia. Fifth edn. Santa Cruz: Asociación Armonia.
- Ridgely, R. S. & Tudor, G. (1989) The birds of South America, 1. Austin: University of Texas Press.
- 3. Schulenberg, T. S., Stotz, D. F., Lane, D. F., O'Neill, J. P. & Parker, T. A. (2007)

 Birds of Peru. Princeton, NJ: Princeton University Press.

Virgilio Yábar Calderón

E-mail: amazonialodge1@yahoo. com.

Received 25 November 2009; final revision accepted 24 February 2010

First record of Red-crested Finch Coryphospingus cucullatus for Manu Biosphere Reserve, Peru

In Peru, Red-crested Finch Coryphospingus cucullatus occurs in dry-land habitats such as in the Marañón Valley, in the north of the country, and the Urubamaba Valley, in dpto. Cusco^{2,3}. However, on 11–13 November 2009, a male Red-crested Finch was present around the garden of Amazonia Lodge (12°52'S 71°22'W) in dpto. Madre de Dios, at 500 m (Fig. 1). The garden is surrounded by mature secondary forest and várzea forest. The bird was usually seen in the company of Silverbeaked Tanagers Ramphocelus carbo or Plain-crowned Spinetails Synallaxis gujanensis. This appears to be the first record for